

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

Applicant(s): Gleave, et al.	
Application No.: 09/913,325	Group Art Unit: 1635
Filed: 8/10/2001	Examiner: Tracy Vivlemore
Title: TRPM-2 Antisense Therapy	Confirmation No: 8469
Attorney Docket No.: UBC.P-020	
Customer No.: 57381	

Commissioner for Patents  
PO Box 1450  
Alexandria, VA 22313-1450

**SUPPLEMENT TO AMENDMENT ACCOMPANYING RCE**

Sir:

Supplemental to the Amendment Accompanying RCE previously filed March 21, 2007, attached is the information regarding PC-3 cells referred to in the amendment.

Respectfully,

Marina Larson & Associates, LLC



Marina T. Larson, Ph.D. Reg. No. 32,038  
P.O. Box 4928  
Dillon, CO 80435  
Tel: (970) 262-1800  
Fax: (970) 262-1809



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search  for

Limits

Preview/Index

History

Clipboard

Details

**Note:** Performing your original search, +*"pc-3" "androgen receptor"*, in PubMed will retrieve 176 citations.

Display  Show  Sort by  Send to

All: 1

Review: 0



☐ 1: [J Steroid Biochem Mol Biol.](#) 2003 Apr;84(5):493-502.

**ELSEVIER**  
**FULL-TEXT ARTICLE**

[Links](#)

**Secretion of endogenous kallikreins 2 and 3 by androgen receptor-transfected PC-3 prostate cancer cells.**

**Kollara A, Diamandis EP, Brown TJ.**

Samuel Lunenfeld Research Institute, Mt. Sinai Hospital, Suite 876, 600 University Avenue, Toronto, Ont., Canada M5G 1X5.

Androgen independent PC-3 cells lack androgen receptor (AR) expression and do not produce kallikrein 2 (hK2) or 3 (prostate-specific antigen, PSA). In this paper, we examined the ability of androgens to stimulate PSA and hK2 production in AR transfected PC-3 cells (PC-3(AR)) and compared this to LNCaP cells. PSA and hK2 were measured in the culture medium and cell lysates using an ELISA-based immunofluorometric assay. Only androgens were able to induce PSA and hK2 secretion in PC-3(AR) cells in a dose- and time-dependent manner depending on the level of AR present. The level of androgen-induced PSA and hK2 secretion in PC-3(AR) cells was approximately 1.5 and 0.9% that induced in LNCaP cells, respectively. Insulin-like growth factor-I (IGF-I), which has been shown to activate AR in the absence of ligand, did not activate PSA secretion in the absence of androgen, but further increased the dihydrotestosterone-induced PSA secretion in PC-3(AR) cells. The lack of PSA and hK2 production in parental PC-3 cells is thus a result of their lack of AR expression. PSA and/or hK2 production in PC-3(AR) cells can thus serve as an endogenous reporter system to investigate AR action or to screen putative endocrine disrupters.

PMID: 12767274 [PubMed - indexed for MEDLINE]

**Related Links**

Dissociation between androgen responsiveness for malignant growth vs. expression of prostate specific differentiation markers PSA, hK2, and PSMA in human prostate cancer models. [Prostate. 2003]

Androgen receptor activation in prostatic tumor cell lines by insulin-like growth factor-I, keratinocyte growth factor, and epidermal growth factor. [Cancer Res. 1994]

Interactive effects of triiodothyronine and androgens on prostate cell growth and gene expression. [Endocrinology. 1999]

Different proportions of various prostate-specific antigen (PSA) and human kallikrein 2 (hK2) forms are present in noninduced and androgen-induced LNCaP cells. [Prostate. 2000]

Tumor necrosis factor-alpha represses androgen sensitivity in the LNCaP prostate cancer cell line. [J Urol. 2000]

[See all Related Articles...](#)

Display  Show  Sort by  Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Mar 6 2007 07:35:03